

## Research Article

# Comparative analysis of open versus laparoscopic ventral hernia repair

Suresh Badiger\*, Sanjay N. Koppad, Ashwin Kulkarni, Harsha Kodliwadmth

Department of General Surgery, SDM College of Medical Sciences, Sattur, Dharwad, Karnataka, India

**Received:** 30 May 2016

**Revised:** 04 June 2016

**Accepted:** 06 June 2016

**\*Correspondence:**

Dr. Suresh Badiger,

E-mail: [drsuresh97@gmail.com](mailto:drsuresh97@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** An incisional hernia develops in 7% to 11% of laparotomy incisions. Laparoscopic repair was applied to the ventral hernias, with the expectation of earlier recovery, fewer postoperative complications, and decreased recurrence rates. This prospective study was performed to compare the outcomes after open and laparoscopic ventral hernia repair.

**Methods:** It is prospective study involved 100 patients with ventral hernia, were subjected either to repair by laparoscopy or to open repair. The open surgical operations were performed by the retro-rectus mesh repair, whereas the laparoscopic repairs were performed using the intra-peritoneal on lay dual mesh (IPOM) repair technique.

**Results:** The mean surgery durations were significantly lower in laparoscopic repair when compared to open repair ( $p=0.000$ ). The mean duration of post-operative analgesics used in laparoscopic group is  $2.84\pm 0.60$  days as compared to open ventral hernia repair  $5.47\pm 2.16$  days ( $p=0.000$ ) which is significant. The mean postoperative stay in hospital was shorter for the laparoscopic group than for the open hernia group (2.66 versus 6.88 days;  $p=0.000$ ). Antibiotics used in laparoscopy group is for  $1.33\pm 0.70$  days as compared to open repair  $2.52\pm 0.98$  days ( $p=0.000$ ). Return to the activity or normal daily work is significantly low in laparoscopic group as compared to open repair of hernia (4.13 versus 13.98 days;  $p=0.000$ ). There were fewer post-operative complications in laparoscopy.

**Conclusions:** The findings demonstrate that laparoscopic ventral hernia repair in our experience was safe and resulted in shorter operative time, fewer complications; shorter hospital stays, and earlier returns to daily activity. Hence, it should be considered as the procedure of choice for ventral hernia repair.

**Keywords:** Incisional hernia, Laparoscopic ventral hernia repair, Open retro-rectus mesh repair, Laparotomy